

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>7</sup> :</b> <b>A61K 35/74, 35/72, A23L 1/03, 1/308</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 00/33854</b> <b>(43) International Publication Date:</b> 15 June 2000 (15.06.00)
<b>(21) International Application Number:</b> PCT/NL99/00755 <b>(22) International Filing Date:</b> 9 December 1999 (09.12.99) <b>(30) Priority Data:</b> 1010770 9 December 1998 (09.12.98) NL <b>(71) Applicant (for all designated States except US):</b> N.V. NUTRI- CIA [NL/NL]; P.O. Box 1, NL-2700 MA Zoetermeer (NL). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> DE JONG, Patricia [NL/NL]; Bolderik 3, NL-6026 AM Maarheeze (NL). VAN LAERE, Katrien [BE/NL]; Kamperfoeliestraat 11, NL-6666 WS Heteren (NL). <b>(74) Agent:</b> DE BRUIJN, Leendert C.; Nederlandsch Octrooibureau, Scheveningseweg 82, P.O. Box 29720, NL-2502 LS The Hague (NL).		<b>(81) Designated States:</b> AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>In English translation (filed in Dutch).</i>
<b>(54) Title:</b> PREPARATION THAT CONTAINS OLIGOSACCHARIDES AND PROBIOTICS  <b>(57) Abstract</b>  The present invention relates to a preparation having a health-promoting action, in particular for the prevention and/or treatment of disorders of the digestive tract, which contains one or more probiotics and one or more non-digestible oligosaccharides. The probiotics are preferably chosen from bacterial strains such as a strain of a <i>Lactobacillus</i> or a <i>Bifidobacterium</i> species and from yeast strains such as a strain of a <i>Saccharomyces</i> species.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

### Preparation that contains oligosaccharides and probiotics

The present invention relates to a preparation which has a health-promoting action, in particular for the prevention and/or treatment of disorders of the digestive tract, more particularly of the intestines.

The application relates in particular to a preparation of this type which contains probiotics and non-digestible oligosaccharides.

It is known that certain microorganisms have both a prophylactic and a therapeutic effect on intestinal diseases, such as intestinal infections. When these microorganisms are administered to humans or animals they will compete with pathogenic bacteria for nutrients and/or adhesion sites on the intestinal wall, as a result of which the number of pathogenic bacteria will decrease and infections are prevented or reduced. Such microorganisms are generally designated by the term "probiotics".

If these microorganisms are to have an optimum action they must reach the intestines alive. A further beneficial effect of the administration of live microorganisms to the intestines is, for example, that they are able to ferment the oligosaccharides present in the intestines, whereby, for example, fatty acids with short chains are formed.

In addition it is advantageous on economic grounds if as many microorganisms as possible reach the intestines alive. With the customary preparations which contain probiotics the percentage of microorganisms that reaches the intestines alive is frequently low.

It is therefore an object of the present invention to provide a preparation that contains such probiotics with which a high percentage of the microorganisms administered reaches the intestines alive.

It is also an object of the present invention to provide a preparation that can be used for the treatment of disorders of the intestines and/or that can be used for prophylactic treatment of the intestines.

The present invention now relates to a preparation having a health-promoting action, in particular for the prevention and/or treatment of disorders of the digestive tract, more particularly of the intestines, which contains probiotics and non-digestible oligosaccharides.

Without wishing to be tied to any theory, it is assumed that the oligosaccharides form a substrate for the probiotics, as a result of which the likelihood that said

microorganisms reach the intestines alive increases and as a result of which the likelihood that they are able, in combination with the oligosaccharides present in the intestines, to exert their beneficial action increases. The oligosaccharides could therefore also be designated as "prebiotics".

5 The probiotics and non-digestible oligosaccharides are present in the preparation in a ratio of 1 to 5 g oligosaccharides per  $10^8$  to  $10^{11}$  cells of the probiotic.

The oligosaccharides used are chosen from the so-called "non-digestible oligosaccharides", that is to say oligosaccharides which are not absorbed by the human or animal body. These oligosaccharides as a rule have a degree of polymerisation of 2 to 20.  
10 This implies that the oligosaccharides consist of 2 to 20 monosaccharide units. In the case of a degree of polymerisation of less than 2, that is to say in the case of the monosaccharide, the preparation is not effective since such monosaccharides are absorbed by the human or animal body. At a degree of polymerisation of greater than 20 the oligosaccharides lose their beneficial action.

15 Preferably the oligosaccharides have a degree of polymerisation of 2 to 10, more preferentially of 2 to 6, more particularly 3 to 10 and more preferentially 3 to 6.

Where reference is made in the present application to oligosaccharides this term is used to refer both to oligosaccharides having one specific chain length and to mixtures of oligosaccharides having different chain lengths. However, a mixture of oligosaccharides  
20 having different chain lengths is preferred.

Furthermore, the oligosaccharides usually do not consist entirely of disaccharides. The disaccharide content is usually less than 90 % and sometimes less than 60 %.

The average degree of polymerisation is as a rule more than 2.1, usually more than 2.5.

25 The oligosaccharides used in the invention are, as a rule, furthermore so chosen that they are at least 20 % usable as substrate for the probiotic microorganisms present in the preparation, as determined by high performance anion exchange chromatography.

Examples of suitable oligosaccharides are transgalacto-oligosaccharides (TOS), fructo-oligosaccharides (FOS) and combinations thereof.

30 It is particularly advantageous if the oligosaccharides are added to the preparation in the form of a hydrolysis product of one or more polysaccharides, for example chosen from  $\beta$ -(arabino)galactans,  $\beta$ -(arabino)xylans,  $\beta$ -glucans,  $\beta$ -glucomannans,  $\beta$ -galactomannans,  $\alpha$ -arabans, inulin and combinations thereof. In addition to the oligosaccharides, such a

hydrolysis product can also contains yet further components, such as monosaccharides and saccharides having a degree of polymerisation higher than 20. However, the hydrolysis product must contain at least 50 % non-digestible oligosaccharides, preferably at least 70 %.

5       The polysaccharides which are preferably hydrolysed are  $\beta$ -(arabino)galactans,  $\beta$ -mannans, and xylans.

It is also possible to use a hydrolysis product of one or more fibres which are mainly made up of the abovementioned oligosaccharides, such as fibres from oats, wheat, potatoes, sugar beet, soya polysaccharides and the like.

10       The hydrolysis of the polysaccharide(s) and/or the fibre(s) can be carried out in a manner known per se, for example by the use of suitable enzymes.

The concentration of oligosaccharides in the preparation is such that 0.5 to 20 gram per day can be administered. If desired, this administration can be spread over the day, as long as the total quantity of the oligosaccharides administered remains in the range  
15       described above. In general the preparation will be administered 2 to 4 times per day.

In general the oligosaccharides will make up 5 wt.% to 50 wt.% of the total preparation.

The probiotics suitable for the present invention are generally known. They comprise, in general, one or more bacterial strains suitable for use in food preparations,  
20       such as lactic acid bacteria suitable for use in food preparations, or one or more yeast strains suitable for use in food preparations, or a combination thereof. These probiotics will usually have GRAS ("Generally Recognised As Safe") status.

Suitable bacterial strains are, for example, chosen from those which are described in European Application 97202900.3 in the name of the Applicant. Other possible probiotics  
25       are the *Pediococci*, *Propionibacteria* or *Leuconostoc* species. The *Lactobacillus*, and *Bifidobacterium* genera and combinations thereof are to be preferred.

The *Bifidobacterium* strain used can be any strain which is suitable for, and preferably is also approved for, administration to humans and animals, such as *Bifidobacterium*, *bifidum*, *Bifidobacterium breve*, *Bifidobacterium lactis* or  
30       *Bifidobacterium longum*, or a combination thereof.

The *Lactobacillus* strain is preferably so chosen that this produces mainly, preferably exclusively, dextrorotatory (L+) lactate. What is meant by this is that the lactate produced is less than 5 %, preferably less than 2 %, laevorotatory lactate. It is, of course, possible

that the microorganism produces other metabolites in addition to lactate and the beneficial action of the microorganism can (also) be based on the formation of these further metabolites.

Examples of these are *Lactobacillus acidophilus*, *Lactobacillus brevis*, *Lactobacillus*  
5 *(para)casei*, *Lactobacillus fermentum*, *Lactobacillus plantarum* and *Lactobacillus rhamnosus*, and combinations thereof.

The yeast strain used in the invention can be any strain which is suitable for, and preferably is also approved for, administration to humans and animals, such as of the genus *Saccharomyces*. Examples of suitable *Saccharomyces* species are *Saccharomyces*  
10 *cerevisiae* and *Saccharomyces boulardii*.

The yeast in the preparation can be alive or dead. Both dead and live yeast contains a high content of mannoproteins, which are able to prevent the adhesion of bacteria to the intestinal wall to a large extent. Administration of dead yeast offers advantages in the case of people who are suffering from inflammatory intestinal diseases.

15 If the yeast is used in dead form, at least one other live microorganism must be used. Said live organism can once again be chosen from both the *Lactobacillus*, *Bifidobacterium*, *Pediococci*, *Propionibacteria* and *Leuconostoc* strains, but can also be live *Saccharomyces*.

It is possible to use both a single microorganism and a mixture of microorganisms.

20 The total concentration of the probiotics is  $10^6$  to  $10^{12}$ , preferably  $10^9$  to  $10^{10}$ , live cells per gram of total product. If a combination of microorganisms is used, the minimum concentration of each of the microorganisms must be such that there can still be said to be live organisms, that is to say at least approximately 10 per gram of product. The total concentration of microorganisms must always be within the above specified range of  $10^6$   
25 to  $10^{12}$  live cells per gram of total product.

If dead *Saccharomyces cerevisiae* is also used, this is administered in a quantity of 0.5 to 5 g per day.

The combination of *Lactobacillus rhamnosus* with transgalacto-oligosaccharide or hydrolysis products of (potato) galactan is found to be a particularly suitable combination  
30 of oligosaccharide and probiotic.

The suitability of a specific oligosaccharide for a specific probiotic can be determined by determining the capacity of the microorganism concerned to ferment said oligosaccharide or said oligosaccharide fraction. A specific method for this is given below

with the examples.

The administration forms of the preparations according to the invention are as a rule analogous to those which have been described in European Application 97202900.3 in the name of the Applicant, the contents of which are incorporated herein by reference. In this context it must be pointed out that according to the present invention also only one probiotic microorganism (including a yeast) can be present and that in addition to the one or more probiotic microorganisms one or more oligosaccharides can also be incorporated, in the quantities specified above.

The preparations according to the present invention can, among others, be administered in the form of a nutritional supplement, total nutrition and clinical nutrition. Reference is likewise made to European Application 97202900.3 for the specific additives which are added to such foods and the preparation and applications of such foods.

The probiotics are preferably added to the preparation in (freeze-)dried form. It is also possible to produce liquid preparations, but these must be stored cool. Furthermore, one or more of the microorganisms can be used in encapsulated form, for example in order to improve the shelf life.

If the preparation according to the invention is used as a nutritional supplement, it can be administered as such, can be mixed with a suitable drinkable liquid, such as water, yoghurt, milk or fruit juice, or can be mixed with solid or liquid food. In this context the nutritional supplement can be in the form of tablets, capsules, powders, sachets, pastilles, sweets, bars and corresponding administration forms, usually in the form of a unit dose.

A supplement according to the invention can, for example, have the following composition:

- probiotics:	10 - 40 wt.%
- oligosaccharides:	40 - 80 wt.%
- further additives:	0 - 40 wt.%,

to a total of 100 wt.%.

The preparation can also be in the form of a food preparation that is suitable for direct consumption, such as total or clinical nutrition. This can be either oral nutrition or nutrition for administration via a tube or catheter.

Such foods can be in solid form or liquid/drinkable form and can contain all customary additives for (total and/or clinical) nutrition, including proteins, vitamins, minerals, trace elements and the like.

A total nutrition according to the invention can, for example, have the following composition:

- probiotics: 0.1 - 10 wt.%

- oligosaccharides: 1 - 20 wt.%

5 - further additives: 75 - 95 wt.%

to a total of 100 wt.%.

According to a particular embodiment, the preparation according to the invention is in the form of a (supplement for a) baby food or a nutritional supplement for babies.

The preparations described above can be used for the same applications as those described in European Application 97202900.3 in the name of the Applicant, in particular in the treatment of disorders of the intestines, such as diarrhoea, such as can arise when travelling or after treatment with antibiotics, or which results from food poisoning. Another application is in the treatment of inflammatory bowel diseases (IBD), such as colitis ulcerosa and Crohn's disease. The preparations according to the invention are also suitable for patients who have a food allergy, such as an allergy to cow's milk or to gluten.

The probiotics and non-digestible oligosaccharides can also be used in baby food to prevent or treat intestinal problems.

The invention will now be explained with the aid of the following non-limiting examples.

20

### Examples

To determine suitable combinations of oligosaccharides and probiotics the microorganisms listed below were tested to determine their capacity to ferment structurally different oligosaccharide fractions. Strains pre-cultured in liquid medium based on thioglycolate (Oxoid, Unipath Ltd, Basingstoke, Hampshire, UK) were subjected to sub-culture in thioglycolate to which 0.5 % (m/V) oligosaccharides were added. The sugar-free thioglycolate medium and the oligosaccharide solutions were sterilised separately for 15 minutes at 121 °C.

Following anaerobic incubation for 48 hours at 37 °C the pH was measured with the aid of a micro-pH meter (Sentron, Roden, The Netherlands). The changes in the residual oligosaccharide content and the formation of reaction products were determined using HPAEC (high performance anionic exchange chromatography). For HPAEC analysis the cultures were centrifuged and the supernatant liquor was diluted 10-fold with H<sub>2</sub>O and



boiled for 5 minutes to stop the enzymatic activity. The purity of the strains was checked before and after fermentation.

The HPAEC system consisted of a Dionex Bio-LC GPM-II quaternary gradient module (Dionex Corporation, Sunnyval, CA, USA) equipped with a Dionex Carbopac PA-100 column (4 \* 250 mm) in combination with a Carbopac PA-100 guard column (3 \* 25 mm). The samples (20 µl) were injected using a Spectra Physics SP8880 autosampler. The oligomers were analysed using a gradient of sodium acetate in 100 mmol.l<sup>-1</sup> NaOH.

The results of these experiments are given below in the table for a number of combinations of probiotics and substrates, where:

- 10    ++        indicates complete fermentation  
       +        indicates partial fermentation  
       -        indicates no or very limited fermentation

Bacteria	Substrate			
	Hydrolysis product of:			
	Arabino- galactans	Arabans	Arabinoxylans	Fructans
Bi. Breve	++	+	-	++
Bi. Longum	++	++	+	++
Bi. adolescentis	++	+	++	++
L. acidophilus	++	-	-	+
L. fermentum	++	-	-	+
K. pneumoniae	++	-	-	+
C. perfringens	-	-	-	++

- 15        A few examples of preparations according to the present invention are given below.

#### Example I: Supplement

- A suspension of *Lactobacillus rhamnosus* ATCC 7469 (Lb) was freeze-dried, a powder being obtained which contained at least 10<sup>9</sup> viable cells per gram powder.
- 20 Transgalacto-oligosaccharides (TOS), obtained from lactose (Borculo Whey Products), were dissolved in water at 40 °C to a solids content of 25 % and this solution was spray-dried. The two powders were mixed in a TOS/Lb ratio of 4/1 until a homogeneous product was obtained. Sachets were filled with 2 - 5 g of this mixture, depending on the dosage

regime (5 g for one sachet per day; 3 g for two sachets per day). The contents of one sachet can, for example, be taken mixed in a glass of orange juice or milk.

**Example II: Synbiotic bar**

5        A 23 g bar was prepared from 4.0 g oat flakes, 4.0 g wheat flakes, 3.0 g puffed rice, 1.0 g crushed hazelnuts, 0.25 g honey, 3.0 g raisins, 1.5 g maltodextrin, 1.0 g freeze-dried *Lb rhamnosus*, 0.5 g baker's yeast (*Saccharomyces cerevisiae*; Gist Brocades) and 5.0 g transgalacto-oligosaccharides.

10      **Example III: Method for hydrolysing vegetable fibres**

      A 20 % suspension of fibres, for example from wheat, potatoes, oats, soya polysaccharides, carob gum or sugar beet, in water was prepared. These sources of fibre are commercially available. The suspension was heated to a temperature of between 20 and 50 °C (preferably 35 - 45 °C), after which enzymes were added in a quantity of one part  
15      enzyme per 5 - 500 parts (m/m) substrate. The choice of the type of enzyme is dependent on the type of polysaccharide. Examples of suitable enzymes are Novoferm Pectinex Ultra s.p.-L, Pentopan and Ultra.s.p. (NOVO).

      After 0.5 - 4 hours the reaction was terminated by heating, after which the solution thus obtained can be used as the oligosaccharide fraction in the preparations according to  
20      the invention, optionally after further filtration/purification or after concentrating.

**Example IV: Synbiotic mixture for mixing with a complete enteral clinical nutrition**

      A mother batch of a powder mixture was prepared in accordance with the method of Example I. The powder consisted of 20 % hydrolysed wheat arabinoxylans, 20 %  
25      hydrolysed potato arabinogalactans, 20 % hydrolysed carob gum, 20 % hydrolysed sugar beet fibre (arabans), 15 % hydrolysed oat fibres (glucans) and 5 % *Bifidobacterium longum*. 5 g of the powder mixture is placed in a sachet. The contents of this sachet can be added to a standard enteral clinical nutrition a maximum of 30 min before use.

30      **Example V: Synbiotic powder mixture for fortifying baby food**

      A synbiotic mixture was prepared in accordance with the method of Example I. The composition contains 10 % baker's yeast (Gist Brocades), 40 % mannoproteins, obtained from yeast, 25 % inulin and 25 % raffinose.

**Example VI: Sweet that contains a synbiotic mixture**

A 2 g sweet was prepared starting from 58 % digestible carbohydrates (glucose syrup), 35 % TOS, 4 % *Lactobacillus rhamnosus* ATCC 7469, and 2 % flavourings and  
5 colourants.

## CLAIMS

1. Preparation having a health-promoting action, in particular for the prevention and/or treatment of disorders of the digestive tract, which contains one or more probiotics and one  
5 or more non-digestible oligosaccharides.
2. Preparation according to Claim 1, wherein the oligosaccharides have a degree of polymerisation of 2 to 20, preferably of 2 to 10, more preferentially of 2 to 6.
- 10 3. Preparation according to Claim 1 or 2, wherein the oligosaccharides have been obtained by the hydrolysis of one or more polysaccharides, chosen from  $\beta$ -(arabino)galactans,  $\beta$ -(arabino)xylans,  $\beta$ -glucans,  $\beta$ -glucomannans,  $\beta$ -galactomannans,  $\alpha$ -arabans, inulin and combinations thereof.
- 15 4. Preparation according to Claim 3, wherein the polysaccharides are chosen from  $\beta$ -(arabino)galactans,  $\beta$ -mannans, xylans and combinations thereof.
5. Preparation according to Claim 1 or 2, wherein the oligosaccharides originate from the hydrolysis of one or more fibres, such as fibres originating from oats, wheat, potatoes,  
20 sugar beet, soya polysaccharides and the like.
6. Preparation according to one of the preceding claims, wherein the probiotics comprise at least one bacterial strain or at least one yeast strain, or a combination of at least one bacterial strain and at least one yeast strain.
- 25 7. Preparation according to one of the preceding claims, wherein at least one of the bacterial strains is chosen from one or more strains of a *Lactobacillus* or a *Bifidobacterium* species and the yeast strain is a strain of a *Saccharomyces* species.
- 30 8. Preparation according to one of the preceding claims which also contains dead yeast cells.
9. Preparation according to one of the preceding claims, wherein the ratio between the

one or more probiotics and the one or more non-digestible oligosaccharides is in the range of 1 to 5 g oligosaccharides per  $10^8$  to  $10^{11}$  cells of the probiotic.

10. Preparation according to one of the preceding claims which contains the probiotics in  
5 a concentration of  $10^7$  to  $10^{11}$  live cells per gram of total product.

# INTERNATIONAL SEARCH REPORT

International Application No.

PCT/NL 99/00755

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K35/72 A61K35/72 A23L1/03 A23L1/308

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K A23L

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 856 259 A (SITIA YOMO SPA) 5 August 1998 (1998-08-05) column 1, line 14-44 column 5, line 32 -column 6, line 12 claims	1-3, 9, 10
P, X	EP 0 904 784 A (NUTRICIA NV) 31 March 1999 (1999-03-31) cited in the application column 1, line 3-5 column 5, line 32-44 column 5, line 54 -column 6, line 23 column 8, line 5-40 claims	1-10

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

21 March 2000

Date of mailing of the international search report

31/03/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3018

Authorized officer

Van Moer, A

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/NL 99/00755

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>WO 96 26732 A (NEWPHARMA SRL ;RONCHI CELESTINO (IT); BERLATI FABIO (IT)) 6 September 1996 (1996-09-06) page 1, line 5 -page 2, line 29 page 3, line 7-11 page 5, line 6-14 claims</p>	1-4,6-10
X	<p>WO 98 26787 A (ARNOTT S BISCUITS LTD ;BURNS PHILP &amp; COMPANY LIMITED (AU); CONWAY) 25 June 1998 (1998-06-25) table I page 4, line 25 -page 5, line 24 claims</p>	1-10
Y	<p>WO 97 35596 A (ABBOTT LAB) 2 October 1997 (1997-10-02) page 3, line 22-25; claims</p>	1,2,5,9, 10
Y	<p>US 5 021 245 A (BORSCHEL MARLENE W ET AL) 4 June 1991 (1991-06-04) claims</p>	1,2,5,9, 10

# INTERNATIONAL SEARCH REPORT

Information on patent family members

Intern. Application No

PCT/NL 99/00755

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0856259 A	05-08-1998	AT 169456 T BR 9706278 A CA 2213113 A CN 1185904 A DE 69600525 D DE 69600525 T JP 10286078 A US 5895648 A	15-08-1998 01-06-1999 23-06-1998 01-07-1998 15-10-1998 10-12-1998 27-10-1998 20-04-1999
EP 0904784 A	31-03-1999	NONE	
WO 9626732 A	06-09-1996	IT MI950378 A	28-08-1996
WO 9826787 A	25-06-1998	AU 5396198 A	15-07-1998
WO 9735596 A	02-10-1997	US 5902578 A	11-05-1999
US 5021245 A	04-06-1991	AT 116110 T AU 640781 B AU 7720991 A CA 2043003 A,C DE 69106271 D DE 69106271 T EP 0471153 A ES 2070365 T GR 3015624 T HK 1006797 A IE 65960 B IL 98137 A JP 2047153 C JP 4228026 A JP 7067368 B NZ 238192 A	15-01-1995 02-09-1993 28-11-1991 23-11-1991 09-02-1995 18-05-1995 19-02-1992 01-06-1995 30-06-1995 19-03-1999 29-11-1995 12-04-1994 25-04-1996 18-08-1992 26-07-1995 25-11-1993